


## Prescribing conscious sedation

- The following applies for prescription of medications used for conscious sedation (e.g. midazolam, fentanyl, morphine)
- A midazolam prescription for 1-5mg given in boluses titrated to effect is shown below as an example
- This prescription should be done prior to beginning a procedure (e.g. OGD) such that the required medications can be administered


### Click order on the black task bar




### Click ENTER



### Type the name (or beginning of it) in the search bar



### Select the route of administration (IV INJ in this case) and click the downward red arrow



### Complete the required information: Dose (1-5mg), Freq (PRN), PRN Reason (sedation), Dose instructions (Titrate to effect)

midazolam [midazolam 5 mg / 5 mL ampoule]

midazolam [midazolam 5 mg / 5 mL ampoule] \*DOSE mg IV INJ \*FREQ SCH 30/10/25 13:00 STOP DATE \* New

midazolam

midazolam infusion IV INF

midazolam [midazolam 5 mg / 5 mL ampoule] 1 - 5 mg IV INJ PRN PRN 30/10/25 14:37 STOP DATE New

\* Dose 1 - 5

Units mg

\* Route IV INJ

\* Freq PRN

\* Sch PRN

Monograph View

Additional Admin Copy Admin


Pending Yes No

\* PRN Reason sedation

Label Comments

\* Dose Instruction Titrate to effect

### Click SUBMIT (top right)



### Click SAVE (top right) and enter your pin

